

# Abstract

J Intern Med. 1989 Oct;226(4):271-6

## Pronounced lowering of serum levels of lipoprotein Lp(a) in hyperlipidaemic subjects treated with nicotinic acid.

Carlson LA, Hamsten A, Asplund A.

King Gustaf V Research Institute, Stockholm, Sweden.

**METHODS:** Thirty-one consecutive unselected hyperlipidaemic patients were treated daily with 4 g of nicotinic acid for 6 weeks. The concentrations in serum of lipoprotein Lp(a), and the major lipoprotein classes, were determined before and after the treatment.

**RESULTS:** Nicotinic acid significantly reduced the serum levels of Lp(a) in the whole patient group. Linear regression analysis showed a strong negative relationship between the percentage reduction of Lp(a) and the serum triglyceride level before treatment ( $r = -0.78$ ), which implied that for patients with a serum triglyceride concentration above 7.5 mmol l<sup>-1</sup> there was a rise of Lp(a). The average individual percentage decrease of the concentration of Lp(a) was calculated after the exclusion of four patients who had serum triglyceride levels above 10 mmol l<sup>-1</sup>. The decrease was 38% with a 95% confidence interval of 28-47%. The absolute decrease of Lp(a) was correlated with the pretreatment levels of Lp(a) ( $r = 0.91$ ). Within the whole group of patients there was a linear relationship between the percentage decrease of Lp(a) and that of LDL cholesterol ( $r = 0.88$ ).

**CONCLUSION:** This latter strong relationship might be due to an inhibition of the synthesis of the protein common to the two lipoproteins, apolipoprotein B.

PMID: 2530298